


### Drug Facts (continued)

#### Inactive ingredients

carbomer 934P, glycerin, methylparaben, octoxynol-9, propylparaben, purified water, sodium hydroxide

Questions? 1-866-255-6960  
or www.fleetlabs.com

Complete enema in a  
disposable squeeze  
bottle with soft,  
pre-lubricated Comfortip®



Fleet® Enemas are **LATEX FREE**. Allergic reaction to materials containing latex is a growing and serious medical problem.

© C.B. Fleet Company, Inc. 2007  
The package design is a trademark of C.B. Fleet Company, Inc.

C.B. FLEET CO., INC. Lynchburg, VA  
24502 U.S.A.

**Fleet**  
Laboratories, Inc.

### How to use this enema:



- shake bottle well before removing protective shield
- **REMOVE ORANGE PROTECTIVE SHIELD FROM ENEMA COMFORTIP® BEFORE INSERTING.**
- with steady pressure, gently insert enema tip into rectum with a slight side-to-side movement, with tip pointing toward navel. Insertion may be easier if person receiving enema bears down, as if having a bowel movement. This helps relax the muscles around the anus.
- do not force the enema tip into rectum as this can cause injury
- squeeze bottle until nearly all liquid is gone. It is not necessary to empty the bottle completely, as it contains more liquid than needed.
- remove Comfortip® from rectum and maintain position until urge to evacuate is strong

### Positions for using this enema:



- **Left-side position:** lie on left side with knee bent, and arms resting comfortably



- **Knee-chest position:** kneel, then lower head and chest forward until left side of face is resting on surface with left arm folded comfortably

# Fleet®



## Enema bisacodyl

stimulant  
laxative  
ready-to-use  
latex free

Soft,  
Flexible  
Comfortip®



1.25 Fl Oz  
(37 mL)

\*Laxative Brand

### Drug Facts

**Active ingredient**  
(in each 30-mL  
delivered dose) **Purpose**

Bisacodyl  
10 mg .....Stimulant laxative/  
bowel cleanser

**Use** for relief of occasional  
constipation or bowel cleansing  
before rectal examinations

### Warnings

For rectal use only

Ask a doctor before using any  
laxative if you have

- abdominal pain, nausea,  
or vomiting
- a sudden change in bowel  
habits lasting more than 2 weeks
- already used a laxative for  
more than 1 week

This product may cause abdominal  
discomfort, faintness, rectal burning,  
and mild cramps

Stop use and ask a doctor  
if you have

- rectal bleeding
  - no bowel movement after use
- These symptoms may indicate a  
serious condition.

Keep out of reach of children. If  
swallowed, get medical help or  
contact a Poison Control Center  
right away.

### Directions

Single daily dosage

adults and children 12 years and over	one bottle
children under 12 years	DO NOT USE

### Other information

- this product generally  
produces a bowel movement  
in 5-20 minutes
- store at temperatures  
not above 86° F (30° C)
- consult **PHYSICIANS' DESK  
REFERENCE®** for complete  
professional labeling
- carton sealed for safety. If seal  
with Fleet® emblem on top or  
bottom flap is broken or  
missing, do not use