

Drug Facts

Active ingredient (in each suppository)

Glycerin 2g Hyperosmotic laxative

Purpose

Uses

- for relief of occasional constipation
- this product usually produces bowel movement in ¼ to 1 hour

Warnings

For rectal use only

Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Ask a doctor before using any laxative if you

- have a sudden change in bowel habits lasting more than 2 weeks
- already used a laxative for more than 1 week

When using this product rectal discomfort or burning sensation may occur.

Stop using this product and consult a doctor if you have

- rectal bleeding
 - no bowel movement within 1 hour of using this product
- These symptoms may indicate a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

Single Daily Dosage (per 24 hours)

adults and children 6 years and over	1 suppository, or as directed by a doctor
children 2 to under 6 years	use Fleet® Pedia-Lax™ Suppositories
children under 2 years	consult a doctor

Insert suppository well up into rectum, to produce laxative action. Suppository is designed to dissolve only partially, which may not be noticeable.

Other information

- store container tightly closed
- keep away from excessive heat
- mouth of jar sealed for safety. If foil imprinted with green Fleet® emblem is missing or torn, do not use.

Inactive ingredients aloe vera leaf, edetate disodium, purified water, sodium hydroxide, stearic acid

Questions? 1-866-255-6960 or www.fleetlabs.com

Mfd. by C.B. Fleet Company, Inc., Lynchburg, VA 24502 USA
©2018 Trade dress is owned by C.B. Fleet Company, Incorporated.
All rights reserved.