

Drug Facts (continued)

Directions

Single Daily Dosage (per 24 hours)

adults and children 6 years and over	1 suppository, or as directed by a doctor
children 2 to under 6 years	use Fleet® Pedia-Lax™ Suppositories
children under 2 years	ask a doctor

Insert suppository well up into rectum.
To produce laxative action suppository is
designed to dissolve only partially, which
may not be noticeable.

Other information

- Store container tightly closed.
- Keep away from excessive heat.
- Mouth of jar sealed for safety. If foil
imprinted with green Fleet® emblem
is missing or torn, do not use.

Inactive ingredients

edetate disodium, purified water,
sodium hydroxide, stearic acid

Questions? 1-866-255-6960



07417

3

Drug Facts

Active ingredient (in each suppository)

Glycerin 2 g.....

Purpose

Hyperosmotic laxative

Use

For relief of occasional constipation

This product usually produces bowel movement in 1/4 to 1 hour.

Warnings

For rectal use only

May cause rectal discomfort or a burning sensation.

Ask a doctor before using any laxative if you have

- abdominal pain, nausea or vomiting
- a sudden change in bowel habits lasting more than 2 weeks
- already used a laxative for more than 1 week

Stop use and consult a doctor if you have

- rectal bleeding
 - no bowel movement within 1 hour of using this product
- These symptoms may indicate a serious condition.

Keep out of reach of children to prevent accidental ingestion. If swallowed, get medical help or contact a Poison Control Center right away.

